

**CRE-LOX MEDIATED
CONDITIONAL REMOVAL OF
MYELOID CELL-DERIVED *Mif*
IN A MOUSE MODEL OF
ORAL CANCER.**

Arham Siddiqui

CRE-LOX MEDIATED CONDITIONAL REMOVAL OF MYELOID CELL-DERIVED *Mif* IN A MOUSE MODEL OF ORAL CANCER.


ARHAM M. SIDDIQUI, NATHAN RYAN, KELVIN ANDERSON, GRETA VOLPEDO, LAUREN MELLEN, STEVE OGHUMU
PHD, ABHAY SATOSKAR MD, PHD

Oral cancer affects people globally with an estimated 657,000 new cases and 330,000 deaths yearly. Macrophage migration inhibitory factor (*Mif*) is a gene that encodes for a cytokine involved in regulating cell-mediated immunity, immunoregulation, and inflammation. Previous research has demonstrated that global *Mif* inhibition results in reduced oral cancer development. This was associated with a decrease in the accumulation of myeloid cell populations in the oral tumor microenvironment. Therefore, our group is interested in elucidating the role of *Mif* expression by myeloid populations in the oral tumor microenvironment. We previously generated floxed *Mif* reporter mice (*Mif*^{F/FI}). Selective breeding between *Mif*^{F/FI} mice and *Cre* knock-in mice driven by the Lysozyme promoter (*Lyz-Cre*) was performed to produce progeny with *Mif* deficiency in myeloid cells. Genotypic characterization for *Mif*^{F/FI} and *Lyz-Cre* was performed by PCR of genomic DNA extracted from mouse ear tissue. Phenotypic characterization of myeloid *Mif* deficiency was performed through flow cytometric analysis on various myeloid cell populations. Selective breeding of the *Mif*^{F/FI} mice with the *Lyz-Cre* mice for three generations yielded myeloid-specific *Mif* deficient mice. Genotypic PCR characterization of *Mif*^{F/FI} mice yielded band sizes of 404 bp while wild-type mice yielded a band size of 221 bp. Furthermore, genotypic characterization of *Lyz-Cre* mice yielded a band size of 700 bp while wild-type mice yielded a band size of 350 bp. Phenotypic flow cytometric analysis of bone marrow-derived macrophages and peritoneal macrophages from myeloid-specific *Mif* deficient mice revealed absence of *Mif* expression in these cells. Furthermore, in these mice, this difference was not observed in lymphoid cells. Our data indicate that, using the Cre-LoxP system and selective breeding conditional *Mif* gene removal in myeloid cells in mice can be achieved to help us gain a better understanding of the role *Mif* plays in the context of oral cancer and other inflammatory diseases.



BACKGROUND

- Oral Cancer, 657,000 new cases and 330,000 deaths yearly¹
- Reduction of inflammation for those with cancer is beneficial
- Macrophage migration inhibitory factor (*Mif*)
 - Regulates cell-mediated immunity, immunoregulation, and inflammation
 - Systemic inhibition leads to better prevention, prognoses and a decrease in tumor promoting myeloid-derived cell populations²
- The Cre recombinase and loxP system
 - LoxP sequence is inserted to flank the *Mif* gene, aka floxed
 - Cre recombinase dimerizes with loxP and removes targeted *Mif* gene




METHODS


- Selective breeding strategy
- Genotypic characterization
 - Ear samples
- Phenotypic characterization
 - Flow Cytometric Analysis
 - Myeloid derived cells
 - Bone Marrow Derived Macrophages (BMDMs)
 - Bone Marrow Derived Dendritic Cells (BMDCs)
 - Peritoneal Macrophages
 - Lymphocytes
 - Splenocytes
 - Lymph Node derived cells

RESULTS – GENOTYPIC CHARACTERIZATION

Lyz-Cre Genotyping




Mif Genotyping




- Examples of each genotype
- Cre expected bands
 - WT 350 bp
 - Lyz-Cre Mutant 700 bp
- Mif Expected bands
 - WT Mif 221 bp
 - Mif fl 404 bp

RESULTS – PHENOTYPIC CHARACTERIZATION

CD4+ Cells



BMDMs



- CD4 cells showed no difference in *Mif* from WT
- Myeloid-derived BMDMs did have a decrease in *Mif*

CONCLUSIONS

- Deletion of *Mif* in myeloid-derived cells through Cre-LoxP
 - Genotypic characterization
 - Phenotypic characterization
- Further analyses

Key References:

¹ "Oral Cancer." World Health Organization, World Health Organization, 12 Sept. 2018, www.who.int/cancer/prevention/diagnosis-screening/oral-cancer/en/.

² Oghumu, Steve, et al. "Deletion of Macrophage Migration Inhibitory Factor Inhibits Murine Oral Carcinogenesis: Potential Role for Chronic pro-Inflammatory Immune Mediators." *International Journal of Cancer*, U.S. National Library of Medicine, 15 Sept. 2016, www.ncbi.nlm.nih.gov/pubmed/27164411.

THANK YOU!